



Ribble Fisheries Consultative Association

Our Ref: RFCA/HARP/RV.001

John Macholc  
planning@ribblevalley.gov.uk  
Council Offices  
Church Walk  
Clitheroe  
Lancashire  
BB7 2RA

Cc:

Jack Spees, Ribble River Trust  
Keith Ashcroft, Environment Agency  
Jeremy Rawkins, Hodder Consultative  
Mark Owen, Angling Trust

**RE: United Utilities HARP Planning Application Application 3/2021/0119**

Dear Sir,

We have been involved in an engagement process with United Utilities as part of the wider HARP Consultation process over the last six months.

After several meetings between local angling clubs and UU, we feel we have exhausted channels of constructive dialogue and feel we must strenuously oppose UU's planning application for the following reasons.

- We have been requesting sight of the Environmental Impact Assessments for these applications since December 2020. To date, these have not been forthcoming.
- We also requested to see tangible evidence of robust mitigation schemes to manage and contain diffuse pollution – particularly associated with the temporary bridges and haul roads at Newton and West Bradford. To date, this has not been forthcoming.

As a consequence, we have sought expert advice from a number of environmental and scientific experts on what a robust pollution monitoring and alert scheme would look like, which is attached to this submission.



In our view, any future EIA produced by UU must contain a monitoring and alert system which is at least as robust as the system outlined below.

Furthermore, before any work commences on the HARP project, this pollution monitoring system must be installed and operating for at least 12 months in order to provide a rigorous set of baseline data.

As the 2020 Fisheries Monitoring report by the Ribble Rivers Trust makes clear, from an environmental perspective, the Ribble is in a precarious state and any increase in the volume of diffuse pollution has the potential to have a catastrophic impact on wildlife.

Projects of the scale and duration of the HARP project will impact the whole spectrum of riparian life on the Ribble and Hodder, affecting not only the globally endangered Atlantic salmon, but also aquatic fly life, birdlife and otters.


The Ribble is now one of only a handful of rivers across England with a viable migratory salmon run – thanks in part to the efforts of local anglers, who voluntarily return more than 90 per cent of migratory salmonids caught on the river.

With an anticipated duration of six to eight years, unless diffuse pollution and sediment run-off are effectively monitored, managed and mitigated, this project has the potential to bring the Atlantic salmon to the point of extinction in the Ribble catchment.

As the representatives of twenty (20) local angling clubs with a combined membership of some Thirteen Thousand Five Hundred (13,500) anglers, we request that these applications are subject to the most robust scrutiny – particularly from the environmental perspective.

We would request the planning authority consult us directly once the EIA is submitted as we have gone to considerable time and expense to produce the attached monitoring proposals in the absence of any tangible monitoring systems proposed by UU.

Yours faithfully



Hon Secretary  
**Ribble Fisheries Consultative Association**



Document	HARP - River Ribble and Hodder Water Quality Monitoring
Customer	Ribble Fisheries Consultative Association
Date	22-06-21

The RFCA as requested by HARP/UU to produce a monitoring framework to oversee the forthcoming project at West Bradford, Clitheroe. Additionally, the Hodder Consultative have previously presented their proposal to HARP/UU.

As with the Hodder Consultative we agree that it imperative that benchmark is established and refer to the historical references detailed within their submission.

The RFCA would expect as a minimum that HARP/UU undertake a commitment to provide a comprehensive monitoring program which would commence between two and one year prior to any works being commenced and would run for at least one year post work.

**The Key Elements would include but not limited to:**

- 1. Ecology surveys** – Long term tracking of ecology trends on the river
- 2. Spot Sampling** - For identifying long term trends and detect elements such as heavy metals.
- 3. Real-Time Monitoring** – To identify events and short-terms changes in the river, also to trigger sample collection or further analysis

We feel that the Hodder Consultative have covered Spot Sampling in respect of Citizen Science using schemes such as the Riverfly Partnership and if required the T&S SmartRivers.

With regards to Real-Time in River Monitoring, we consulted with Matt Loewental (Environment Agency Lead Water Quality Instrumentation). Matt Loewental recommend that we engage Meteor Communications to provide an outline of which they have supplied below.



**Scope:**

To provide real time Water Quality monitoring to support a wider programme of monitoring on the River Ribble and River Hodder to assess the impact of the HARP works. Monitors would be located U/S and D/S of the proposed working areas. 30 minute real time data with following indicative measured parameters:

- Temperature
- Conductivity
- pH
- Dissolved Oxygen (Optical)
- Turbidity

Alarms can be set on the above parameters to provide early warning of potential events, sent via email or SMS.



## Proposed Equipment:

It is proposed that all sites would have standardised monitoring equipment consisting of the following elements:



### ESNET2-L Portable Monitoring System

Complete portable station in IP66 Enclosure for direct deployment in riverside locations. Inputs for multiparameter sondes, level transducer, additional instrumentation as required.

Real time data via 3G/GPRS



### Xylem EXO2 Multiparameter Sonde

Multiparameter instrument equipped with Temperature, Conductivity, pH, Turbidity and Dissolved Oxygen probes. Complete with central wiper and anti-fouling guard.

Instrument to be deployed directly into water course



### 20W Solar panel kit

Sufficient for 30 min data – 30W may be required for shaded areas

The equipment specified is used by the Environment Agency, Thames Water, Wessex Water, Yorkshire Water, Southern Water, Anglian Water and many consultants and other public bodies for provision of high resolution, real time monitoring.

Similar systems have been and are currently in use on a number of other watercourses in the United Utilities catchment including the River Ehen and River Leven.

### Lease system – Complete Managed Service

This option gives a high degree of flexibility in terms of sensor provision.

It is proposed in this case that the field work would be conducted by the Ribble Fisheries Consultative Association with calibrations at Meteor laboratories, so specification would entail:

- Equipment (ESNET, EXO sonde, Solar panel, connection cable)
- Measured Parameters – Temp, Cond, DO (Optical), Turbidity, pH
- 30 minute Real time data
- MeteorCloud web portal data access
- Calibration every 2 months at Meteor Lab
- Calibration records – Report for each instrument calibration available and electronically stored
- Courier collection and delivery of sondes



**NOTE-** 4 units would be required to provide monitoring at 2 U/S and 2 D/S locations

For purposes of comparison it is possible to provide a basic monitor which gives a broad view of river health by monitoring Temperature, Dissolved Oxygen and Conductivity. This requires basic monthly maintenance and occasional calibration check which can be performed in the field. Equipment service would be as follows:

- Lease of equipment (ESNET, Sonde, Solar panel, connection cable)
- Measured parameters – Temp, Conductivity, Dissolved Oxygen (Optical)
- 30 minute Real time data
- MeteorCloud web portal data access

All field work and calibration is performed by Ribble Fisheries Consultative Association

### **Quality Assurance**


One of the key components of the Meteor service is to monitor the Water Quality data in a proactive and holistic way.

This is achieved through examination of the current data and previous trends for every site, each working day. This enables the team to build a picture of the Water Quality characteristics of the catchments. This is reinforced through automated alarms which also flag suspect data.

### **7.0 Data Representation and Sharing**

The MeteorCloud portal is the primary tool for data dissemination and display. This enables users to view data in graphical, tabular and geospatial modes.

MeteorCloud provides a clear and intuitive interface to view complex data. A typical example display:






This shows a typical rain event passing through a catchment, specifically:

- Conductivity (Blue) – drop as rain dilutes the river water
- Turbidity (Brown) – increase as runoff washed into the river
- Temperature (Red) – Drops as a result of the rain
- Dissolved Oxygen (Light Blue) – Slight increase during the event

The MeteorCloud portal is used across the EA and other Water utilities and is a familiar tool for many Water Quality data users. As part of the ongoing service the following elements are included:

- User training – Online training in use of the MeteorCloud portal
- Telephone and email support
- User setup – Setup of new users and security levels
- Alarm setup – Basic alarm setup and maintenance

### **West Bradford – River Ribble**



***Upstream monitoring point:***



It is proposed to use the structure of the old pumping station building to suspend the sonde into the river. This is sufficiently far upstream of the proposed works to provide a consistent baseline measurement of the river:



Baseline Measurements – Taken at 09.50am on 15-06-2021:

Parameter	Measurement
Temperature	16.6° C
Conductivity	396 $\mu$ S
Turbidity	3.4 NTU
Dissolved Oxygen	120%

All parameters are within typical ranges for a healthy river in the geographic area

***Downstream monitoring point:***

It is proposed to locate the equipment on the weir structure:



This is sufficiently downstream of the works to encompass all of the working area, allowing for the topography of the area and tributaries which may be impacted.




Baseline Measurements – Taken at 10.30am on 15-06-2021:

Parameter	Measurement
Temperature	17.34° C
Conductivity	437 $\mu$ S
Turbidity	4.0 NTU
Dissolved Oxygen	152.8%

All parameters are within typical ranges for a healthy river in the geographic area. The data shows a small variance due to potential impact from tributary and river warming during the day.

### **Marl Hill – River Hodder**





***Upstream monitoring point:***

It is proposed mount the instrumentation onto the existing stone bank wall with the sonde suspended into the river. This is sufficiently far upstream of the proposed works to provide a consistent baseline measurement of the river:




Baseline Measurements – Taken at 11.50am on 15-06-2021:

Parameter	Measurement
Temperature	14.1° C
Conductivity	91 $\mu$ S
Turbidity	1.8 NTU
Dissolved Oxygen	105%

All parameters are within typical ranges for a healthy river in the geographic area.

***Downstream monitoring point:***

It is proposed to mount the downstream monitor on the pipeline bridge structure downstream of the Gibbs Tributary which may also be impacted by the works:



Baseline Measurements – Taken at 12.30am on 15-06-2021:

Parameter	Measurement
Temperature	13.9° C
Conductivity	182 $\mu$ S
Turbidity	0.7 NTU
Dissolved Oxygen	106%

All parameters are within typical ranges for a healthy river in the geographic area.


### **Summary**

The proposed systems for these sites would provide a clear indication of baseline river conditions upstream and downstream of the intended working areas.

Used in conjunction with spot sampling and ecological surveys this will provide a comprehensive view of any potential impact of the works on the river.

It is recommended that the systems are deployed 12 months before the start of any works to give a complete and representative baseline of the river conditions across the four seasons.

Alarms can be set against the measured parameters to enable further sampling to take place or trigger autosampler runs.



# FISHERIES MONITORING OF THE RIBBLE CATCHMENT 2020

The Ribble Rivers Trust

## ABSTRACT

2020 marks the 13th year of the Ribble Trusts inter-annual fisheries monitoring programme. Results and observations from this work helps inform the trust on the productivity of sub-catchment fisheries and where to direct conservation efforts.

Adam Wheeler  
Fisheries Monitoring Officer

# Fisheries Monitoring of the Ribble Catchment

Ribble Rivers Trust  
C/O Hanson Cement  
Ribblesdale Works  
Clitheroe  
BB7 4QF

Phone: 01200 444452

E-mail: [admin@ribbletrust.com](mailto:admin@ribbletrust.com)

**Report title:** Fisheries Monitoring of The Ribble Catchment 2019

**Report reference:** RRT\_ Electric\_Fishing \_2020\_Report

**Report version:** 1.0

**Date:** 01/11/2020

**Prepared for:** Ribble Rivers Trust

**Authored by:** Adam Wheeler - Fisheries and Projects Officer

**Checked by:** Mike Forty – Head of River Conservation

Copyright Ribble Rivers Trust, 2020.

This report has been prepared using due skill, care and diligence for the exclusive use of the commissioning party by Ribble Rivers Trust. No liability is accepted by Ribble Rivers Trust for the use and or application of the contents of the report.

# Fisheries Monitoring of the Ribble Catchment

## Table of Contents

Acknowledgments .....	i
Executive Summary .....	1
1.0 Introduction.....	4
1.1 Sub-Catchment Map.....	5
2.0 Methodologies.....	6
2.1 Electric fishing Surveys .....	6
3.0 Monitoring Results .....	8
3.1 Brown Trout ( <i>Salmo trutta</i> ).....	8
3.2 Atlantic Salmon ( <i>Salmo salar</i> ) .....	11
3.3 Other Species.....	13
5.0 Discussion .....	14
6.0 References.....	18
8.0 Appendices .....	20
8.1 Appendix A .....	20
8.2 Appendix B.....	21
8.3 Appendix C.....	25

# Fisheries Monitoring of the Ribble Catchment

## Acknowledgments

The core fisheries monitoring of the Ribble catchment has been supported and funded by Ribble Rivers Trust and its members. Funding has also been issued from the European Regional Development Fund as part of the European Structural and Investment Fund's Growth Programme 2014-2020, for the monitoring of the Primrose Lodge Blue and Greenway project.

The Ribble Trust is grateful to all involved landowners for their permission to land-access and look forward to continued cooperation for future work. In addition, RRT wishes to thank the staff for their continued hard work and contributions to this year's fisheries programme, which has proved invaluable during this difficult year.

A thanks to all the volunteers that would normally contribute to our survey efforts. We look forward to welcoming you back, hopefully next year.

# Fisheries Monitoring of the Ribble Catchment

## Executive Summary

Despite 2020 being a very difficult and unprecedented year the Ribble Rivers Trust (RRT) concluded its 13<sup>th</sup> year of electric fishing surveys on the Calder, Hodder and Ribble catchments. Whilst the COVID-19 pandemic limited working, strict government guidance was observed to carry out surveys, and additional protective measures were implemented by Ribble Rivers Trust to guarantee the health, safety and wellbeing of their staff. The start of the survey season was also delayed until July 7<sup>th</sup> when restrictions were reduced, and it was deemed safe to proceed. To ensure that all survey work was completed before Environment Agency consents to survey expired, sites were prioritised by their long-term data series and sites that have higher conservational importance. Overall, 198 sites were surveyed.

The methodologies applied to the Ribble Rivers Trust fisheries programme are adapted from Crozier and Kennedy's – 'Application of Semi-quantitative Electrofishing to Juvenile Salmonid Stock Surveys' (1994) and Zippin's – 'Removal Method of Population Estimation' (1958). National Fisheries Classification Scheme (NFCS) grades are then used as a metric for the standardisation of results which are comparable to national datasets. Outputs are used to support and identify future works on the catchment as well as monitoring the long-term impacts of river restoration schemes.

With anthropogenically driven climate change, we are seeing more extreme weather patterns, increased probability of high intensity rainfall events and longer dry periods. This winter (December, January, February) has been the 5<sup>th</sup> wettest winter on record (data back to 1862) for the UK as a whole and some of the highest river levels recorded since December 2015 and March 2019 (Figure i). Following on from this 2020 has also given us the driest May on record with just 17% of the average rainfall. With observation on long term trends, these extreme weather events in addition to unresolved habitat and water quality issues, are having a cumulative effect on the Ribble's salmonid (Atlantic salmon and brown trout) productivity




Figure i. February and May 2020 Met office rainfall data as % of 1981 – 2010 average [www.metoffice.gov.uk](http://www.metoffice.gov.uk)

## Fisheries Monitoring of the Ribble Catchment





Figure ii. Cumulative densities of Atlantic salmon fry recorded on the Calder (50 sites), Ribble (44 sites) and Hodder (44 sites) sub-catchments from 2010 - 2020.

Fish communities have evolved to live in dynamic river habitats and populations have been adaptable to temporal variation. However, the increasing rate of environmental change is far exceeding the capacity of many populations to adapt to new conditions. In particular, cold-water freshwater fish species are predicted to experience strong selective pressures from a wide range of interacting human stressors as well as severe hydrological events and the impacts of warming river habitat. A stark comparison of long-term sites show that this year's Atlantic salmon fry are at their lowest recorded densities on the Ribble and Hodder with less than 100 fry per 100m<sup>2</sup> recorded over the 44 long term sites on each sub-catchment (figure ii). Degradation of Atlantic salmon populations at this rate may lead to unrecoverable numbers by the end of the next decade with local extinction on the Ribble and its sub-catchments.

While normally the catchment reports A and B grade sites, this is the first year where there have been no A or B grades found. Further, this year only one site has been classified as Fair (C-grade), a site is on the main stem Hodder in a location that regularly returns 100s of fry per 100m<sup>2</sup>. The remaining sites on the catchment have all returned D grade or below indicating poor to very-poor fry densities. From the 198 sites surveyed in 2020 the distribution of salmon fry across the catchment is at 30% of the known distribution recorded between 2008 and 2020 by the Ribble Trust. Salmon have a highly complex life cycle that is impacted at every stage by resource competition, environmental stochasticity, and anthropogenic pressures. There is no golden bullet to solve these issues and to have a chance of improving stock of salmon on the Ribble catchment, more needs to be done to reverse negative effects. With such low densities of salmon and trout fry recorded in 2019 and 2020 community composition is being altered and there are concerns to be had over how these cohorts will affect the future adult spawning population. The effective management of freshwater habitat is essential to maintain and enhance salmon and sea trout populations, particularly as there are more unknown and uncontrolled impacts once they reach the saltwater feeding stage of their lifecycle.

## Fisheries Monitoring of the Ribble Catchment



*Figure iii. Cumulative densities of brown trout fry recorded on the Calder (50 sites), Ribble (44 sites) and Hodder (44 sites) sub-catchments from 2010 - 2020.*

Brown trout on the catchment have also struggled with egg to fry survival with 2019 and 2020 producing some of the worst densities recorded by the trust in 13 years (Figure iii). From data collected and seasonal observation it is thought that high flows between spawning and fry emergence have had a marked effect on 0+ salmonid densities. Spate events are important for cleaning gravels and allow the migration of fish to spawning ground. However, extreme events, especially at critical lifecycle stages, can have a large impact on spawning success and the survival of egg to fry. With lower fry densities recorded in 2019-2020, it is hoped that with more resource availability and reduced competition that these cohorts will have a greater chance of surviving to maturity.

# Fisheries Monitoring of the Ribble Catchment

## 1.0 Introduction

The Ribble Rivers Trust (RRT) has been conducting habitat restoration schemes and facilitating improved land management for over twenty years in the Ribble Catchment. With more than 6,000 miles of watercourses the ultimate aim of the RRT is to preserve a resilient system in which there will be resources and habitat to support and sustain fish and wider animal populations and increase biodiversity.

Principally, our continuing aims are to: -

1. Assess the overall status of the juvenile population of salmonids.
2. Monitor the inter-annual variations of the salmonid young of year population.
3. Determine underperforming areas and direct improvement works.
4. Capture the effectiveness of previous habitat improvement works.
5. Generate data and evidence in support of and to report on grant bids and applications.
6. Generate knowledge of rare species to inform responsible development.
7. Locate ecological threats posed by invasive species.
8. Derive future research questions

Despite 2020 being a very difficult and unprecedented year The Ribble Rivers Trust (RRT) concluded its 13th year of electric fishing surveys on the Calder, Hodder and Ribble (Figure 1.1). Whilst the COVID-19 pandemic did limit working, a reduced programme was planned and 195 sites were surveyed with priority given to those with a long-term data series or where considered to have higher conservational importance.



# Fisheries Monitoring of the Ribble Catchment

## 2.0 Methodologies

### 2.1 Electric fishing Surveys

The Trusts applied methodology is adapted from Crozier and Kennedy (1994) and has been in operation since 2008. Riffle/pool habitat is targeted to capture both the young of year and the +1 populous of salmonids using an electric fishing backpack system. Two types of survey are undertaken: semi-quantitative, where the river is actively fished for five minutes covering a measured un-isolated area; and quantitative, where a demarcated area of river is sampled over sequential depletive passes. Salmonid fork lengths are recorded in millimeters at each site and the abundance of other species is noted.

For 2020, 220 survey sites were identified for assessment with priority given to sites that hold the most significant data set, with eight or more years of continuous data or in key locations for monitoring restoration works. Additional sites were selected on the River Brennand, Dunsop and Whitendale as a conservationally important area for salmon spawning. Due to COVID restrictions and delays the survey team maned 35 days field work between 7th July until 30th of September, with a total of 195 sites were completed.

From the above activities the young of year are determined by establishing a maximum fork length discerned from the frequency-length distribution of the species. This method is applied to each major catchment individually to reflect the temporal and spatial differences in fry as the electric fishing season progresses (Appendix B.1 – B.7). Quantitative surveys provide fry densities per 100m<sup>2</sup> from the depletion of a known measured area, these densities are generated from Zippin's (1956,1958) K-Pass Removal method using the FSA package in R version 3.1.0 (R core Team, 2019), whereas, semi-quantitative results are calculated from the number of fry captured in an active five minutes. The equation applied to the semi-quantitative results is formed from the quantitative fry population relationship between a 5 minutes fry capture in the first pass and the total electric fishing result (fry per 100m<sup>2</sup>) (Appendix C.1 and C.2). Data used must reflect the variation in fishing results based on the constant effort of the electric fishing team for each site surveyed. This relationship uses quantitative data collected as well as the addition of a zero, zero point to represent a total absence of salmonids. The resulting equation is taken from the fitted linear regression for 0 + salmonids where:

$$\ln(y + 1) = a + b \ln(x + 1)$$

## Fisheries Monitoring of the Ribble Catchment

The densities of trout and salmon fry per 100m<sup>2</sup> are allocated a grade-score (Table 2.1) which standardises the Trust's field observations with those of the National Fisheries Classification System (NFCS).

Grade	Description	Trout fry per 100m <sup>2</sup>	Salmon fry per 100m <sup>2</sup>
A	Excellent	>38	>86
B	Good	17 – 38	45 – 86
C	Fair	8 – 16	23 – 44
D	Poor	3 – 7	9 – 22
E	Very Poor	1 – 2	1 – 8
F	No Fish Present	0	0

Graded results are transferred to a map layer using ArcGIS 10.3.1 to display catchment scale results. Within the result section the inter-annual comparison of data is based on sites which hold 11 years of consecutive data and the grade change evaluation is the comparison of all sites fished in both 2019 and 2020. Grade results have been organised within the analysis of this report according to geographical coverage determined by sub-catchment.

Maps incorporate the following data files under copyright: © Environment Agency copyright and / or database rights 2020. All rights reserved; © Natural England copyright. Contains Ordnance Survey data © Crown copyright and database right 2020. Base-map imagery sources: National Geographic, Esri, DeLorme, HERE, UNEP-WCMC, USGS, NASA, ESA, METI, NRCAN, GEBCO, NOAA, increment P Corp. All maps © 2020, Ribble Catchment Conservation Trust.


All images © 2020, Ribble Rivers Trust.

## 3.0 Monitoring Results

### 3.1 Brown Trout (*Salmo trutta*)

Brown trout, as an indicator species, can provide the best insight into a catchment's health. This is due to the majority of the population spending their complete lifecycle residing within the river system. In the past fluctuations in brown trout numbers have been observed, however, there has been a noted decline in the densities of fry captured in surveys (Figure 3.1.1). Previous reports have focused on the cumulative grade score of the data series in order to highlight long-term trends, however the loss of salmonids in the high-grade classes are not always accounted for. For example, a site can be classed as an A grade if it has more than 38 fry/100m<sup>2</sup>, yet within the catchment there are sites that can produce fry densities in excess of several hundred. This means that a site can still retain its A grade status despite a high percentage loss in productivity, providing it remains above 38 fry/100m<sup>2</sup> threshold.

From the 11-year dataset, calculating an average at specific intervals set smooths out the data by reducing the impact of fluctuations in short term data. This makes it easier to see overall trends, and with a five-year moving average this downward trend in fry densities is concerning. Given the average lifespan of a river trout is 5 years, then a 5 year moving average gives an indication of the potential health of the entire population, not just the spawning population




*Figure 3.1.1. Cumulative brown trout fry densities on the Ribble catchment for 138 sites fish 2010 – 2020 including 5 year moving average.*

One major impact on fry densities is thought to be the cumulative effect of extreme flow events at critical lifecycle stages. With such low densities of trout fry, there are concerns to be had over how 2019 and 2020 fry populations will affect the future adult populations on the catchment; especially when mortality rates in the first year of life are typically approaching 90% or greater. However, with lower fry densities recorded in 2019-2020, resource availability and reduced competition may allow for these cohorts to have a greater chance of surviving to maturity as survivorship in subsequent years is greatly improved with mortality rates between 40-60% (Jensen, et al., 2008).

# Fisheries Monitoring of the Ribble Catchment

## The Ribble Rivers Trust: Fisheries Monitoring Programme



© The Ribble Rivers Trust. All rights reserved. This document is the property of The Ribble Rivers Trust. It is not to be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage and retrieval system, without the prior written permission of The Ribble Rivers Trust. The Ribble Rivers Trust is not responsible for any loss or damage arising from the use of this document.

**Figure 3.1.2** Catchment map [1:250,000] showing brown trout fry NFCS grades from 195 surveys undertaken by RRT in 2020. Green to red points indicate higher to low grades. Black indicates an absence of trout fry.

## Fisheries Monitoring of the Ribble Catchment

With all sub-catchments underperforming when compared to historical data, sites that would be expected to produce fry densities in the C to A grade bracket have returned poor fry numbers, such as Wigglesworth Beck, Swanside Beck, the lower reaches of Ings Beck on the Ribble catchment and others in the upper reaches of the River Calder. For instance, sites near the confluence of Ings and Swanside Becks have historically produced excellent numbers of both salmon and brown trout fry with an average of 70 trout fry/100m<sup>2</sup> and 79 salmon fry/100m<sup>2</sup>. However, in 2020 only 1.99 trout fry/100m<sup>2</sup> were recorded and no salmon fry were present. Sites that have returned consistently good results appear to be in the upper reaches of the smaller tributaries, where impacts from high spate events are not normally observed.

On a more positive note, fresh run sea trout were recorded on a small spawning tributary of the Hodder catchment after spates in August. These fish can be responsible for a large proportion of young, especially when larger females can produce 1,000-2,000 eggs/kg of bodyweight. Anadromous trout eggs are normally larger than resident brown trout and offspring emerged on average earlier and at a larger body size than the resident fish. This gives fitness benefits to juveniles as larger fry have greater survival, can feed on a wider range of prey, and have better swimming ability.



*Figure 3.1.3 Fresh run sea trout captured in surveys on small spawning tributary of the Hodder Catchment*

# Fisheries Monitoring of the Ribble Catchment

## 3.2 Atlantic Salmon (*Salmo salar*)

The Ribble, once considered a premium English salmon water, continues to see a downward trend in its fry population. In addition to a narrowing distribution of spawning sites, waterbodies that have previously been classed as 'strongholds' for salmon spawning are also being impacted and returning poor fry numbers. These areas must be protected and an increase in conservational efforts must be applied to improve habitat and water quality for the long-term sustainability of this species. Degradation of Atlantic salmon populations at this rate may lead to unrecoverable numbers by the end of the next decade with local extinction on the Ribble and its sub-catchments. There are many issues that are impacting our native fish and without drastic change they will continue to struggle to fulfil their basic biological needs let alone complex lifecycles.

11-year dataset (138 sites): Atlantic salmon




Figure 3.2.1. Total densities of salmon fry for sub-catchments electric fishing sites with 10 years of consecutive data for Atlantic salmon 2010 to 2018.

While normally the catchment reports A and B grade sites, this is the first year where there have been no A or B grades found. Further, this year only one site has been classified as Fair (C-grade), a site on the main stem Hodder in a location that regularly returns 100s of fry per 100m<sup>2</sup>. The remaining sites on the catchment have all returned D grade or below indicating poor to very-poor fry densities (Figure 3.2.2). From the 198 sites surveyed in 2020 the distribution of salmon fry across the catchment is at 30% of the known distribution recorded between 2008 and 2020 by the Ribble Trust (Figure 3.2.2 & 3.2.3).

The Ribble Rivers Trust: Fisheries Monitoring Programme




Figure 3.2.2. Catchment map [1:250,000] showing Atlantic salmon fry NFCS grades from 195 surveys undertaken by RRT in 2020. Green – red points indicate higher to low grades. Black indicates an absence of salmon fry.

The Ribble Rivers Trust: Fisheries Monitoring Programme




Figure 3.2.3. Catchment map [1:250,000] showing Atlantic salmon fry locations historically recorded by RRT between 2008 and 2020.

# Fisheries Monitoring of the Ribble Catchment

## 3.3 Other Species

Bullhead (*Cottus gobio*) remain the dominant non-targeted species on the catchment found within 86.7% of sites (Figure 3.3.1 and 3.3.2). Stone loach (*Barbatula barbatula*) have been recorded in 51.8% of sites and common minnow (*Phoxinus phoxinus*) in 32.8% of sites. The number of European eel (*Anguilla anguilla*) caught as by-catch by the Ribble Trust has seen an annual decrease over the past six years. As a non-targeted species this reduction does not truly reflect the population, but as this species is classed as critically endangered, the reduction is worth noting.

One observation that was made during this electric fishing season was that stone loach and bullhead were noticeably smaller than in previous years and the more mature adults were lacking within sample sites.




Figure 3.3.1. Dominant by-catch by species % presence of sites from 2016 to 2020.




Figure 3.3.2. Accompanying by-catch by species % presence of sites from 2016 to 2020.

## 4.0 Discussion

Fish communities have evolved to live in dynamic river habitats and populations have been adaptable to temporal variation. However, the increasing rate of environmental change is far exceeding the capacity of many populations to adapt to new conditions. In particular, cold-water freshwater fish species are predicted to experience strong selective pressures from a wide range of interacting human stressors as well as severe hydrological events and the impacts of warming river habitat.

A growing body of evidence suggests that high flows between spawning and fry emergence have a marked effect on 0+ salmonid densities (Gillson, et al., 2020), which may be positive or negative. Salmonids use spate events to migrate upstream to spawning grounds, however, beyond an optimum discharge fish migration becomes impeded. Pre-spawning high flow events can provide clean gravel free of organic sediments for spawning, improving oxygen levels and egg survival. After spawning, riverbed scour or fine sediment loading can increase egg mortalities, eggs can also be displaced away from optimal habitat for development. Timing of these events can also be critical to different species; high discharges might increase spawning site availability for salmon and decrease egg-to-fry survival for trout.

Extremes, particularly high flow events between spawning and fry emergence, are thought to be impacting the Ribble catchment. Summer 2018's lowest flows coincided with the main run of sea trout in July and in the Autumn months minimal spate events took place from the end of September delaying returning Atlantic salmon. In 2019 rainfall was exceptionally high for the North West in March with Storms Freya, Gareth and Hannah bringing bands of heavy rain fall and high winds (Figure 4.0.1). This resulted in flood warnings being issued and some of the highest river levels recorded since the winter of 2015. By the month-end, half of the hydrological areas in North West England had observed around twice the expected monthly rainfall for March at a time where fry are emerging for their first feed. In 2020 three named storms crossed the UK during February, Ciara, Dennis and Jorge. The heavy rainfall throughout the month resulted in some severe impacts with many areas flooded and the gauging station on the River Ribble at New Jumbles Rock (among other) logging the highest level on record (Figure 4.0.2). At this time of year salmonid fry are still relying on yolk reserves, remaining very immobile within redds. With river levels exceeding all previous records, heavy sediment movement will have washed out redds leading to high mortalities of undeveloped young that have little to poor swimming ability.




Figure 4.0.1. February May 2020 Met office rainfall data as % of 1981 – 2010 average [www.metoffice.gov.uk](http://www.metoffice.gov.uk)

## Fisheries Monitoring of the Ribble Catchment





Figure 4.0.2. River level data from River Ribble at New Jumbles Rock 2019 and available data for 2020.

For Atlantic salmon we are seeing a year on year decline in recruitment. Salmon have a highly complex life cycle that is impacted at every stage by resource competition, environmental stochasticity and anthropogenic pressures. There is no golden bullet to solve these issues and to have a chance of improving stock of salmon on the Ribble catchment, more needs to be done to reverse negative effects. With such low densities of salmon and trout fry recorded in 2019 and 2020 community composition is being altered and there are concerns to be had over how these cohorts will affect the future adult spawning population. The effective management of freshwater habitat is essential to maintain and enhance salmon and sea trout populations, particularly as there are more unknown and uncontrolled impacts once they reach the saltwater feeding stage of their lifecycle.

One issue impacting salmon stocks is that of man-made barriers on main stem rivers. Atlantic salmon (*Salmo salar*), known as 'the leaper', are renowned for their ability to clear obstacles during their spawning migration. However, a number of studies have documented the movement of Atlantic salmon to key spawning areas may be delayed up to several weeks by man-made obstacles such as weirs, natural waterfalls and even fish passes (Thorstad, et al., 2008). Newton et al. (2018) recorded mean delays of 47.8h at a low-head weir with delays ranging from 15 min to 31 days. Fish may even abandon their migration and enter neighboring watercourses due to the abundance or size of in-river barriers (Thorstad, et al., 2008). The Ribble Trust and EA spring salmon tracking study (2012-2014) showed delays at Arnford weir of between 1 to 24 days. Out of the 11 salmon tagged, 6 were unsuccessful in migrating above this barrier with a maximum of 22 failed attempts to ascend from one individual. For the salmon that were successful there was still an average delay of 3 days at the structure. The sustained and prolonged swimming performance of Atlantic salmon reduces over time, losing between 50 and 70% of their

## Fisheries Monitoring of the Ribble Catchment

total energy content during river migration and spawning (Booth, 1998; Jonsson, et al., 1997). Prolonged delays will prevent salmon from reaching suitable resting or spawning areas and as a consequence will reduce their reproductive success.



*Figure 4.0.3. Partial weir removal on the River Ribble adjacent to Settle sewage works*


Many of the structures on the catchment have associated fish passes which aids upstream migration. However, fish passes rarely achieve 100% efficiency and ascent success decreases with increased slope (Dodd, et al., 2017). Another major consideration is that downstream migrants can also be impacted by weirs and their upstream impounded reaches as Salmon smolts, when encountering a velocity gradient such as accelerating flow over a weir, causing delays to their migration in comparison with open stream sections. Aarestrup and Koed (2003) showed an average delay of 7 days for smolts negotiating weirs with a 0.6 - 2.5 m head drop during emigration. 53% mortality rates were recorded in Atlantic salmon smolts and 18 to 71% documented in brown trout. Delays can result in desmoltification due to a narrow migration window in which to reach the estuary (Leaniz, 2008). These effects are further compounded by unfavorable climatic conditions resulting in low flows during the smolt migration (Gauld et. al., 2013).



*Figure 4.0.4. Partial weir removal on the river Ribble in the Upper Ribble near Horton-in-Ribblesdale*

## Fisheries Monitoring of the Ribble Catchment

Weirs are recognised as predator hotspots and can facilitate excessive predation in a number of ways. Firstly, they offer a vantage point for predators as migrants are forced to swim over shallow crests which are exposed in lower flows. Leaniz (2008) indicated the distribution of cormorants (*Phalacrocorax carbo*) along river courses can be closely related to the location of weirs as these structures cause crowding of fish downstream of the structure. Also, otters (*Lutra lutra*) and cormorants can use the homogenous flow in the upstream reaches to their advantage as prey can be targeted with less energy expenditure. Weir removal will reinstate pool/riffle features, reducing predation success though increased habitat complexity, as variation in flow and depth allows for increased prey evasion. Overcrowding of fish downstream of impoundments have also been shown to facilitate the spread of parasites and infectious diseases, magnifying the impact of pollution incidents and increasing the risk of mass mortalities in low flows (Leaniz, 2008).



*Figure 4.0.5. Samlesbury Weir removal on the River Ribble, Samlesbury. Today, the widest weir removal in England*

This year the trust has completed 3 weir removals including one of the widest in England, bringing the total to 17. More structures need to be offered up for full or partial removal. Our catchment is suffering from the effect of anthropogenic pressures and climate change. The Ribble Trust will continue to improve the water environments of the catchment, by restoring and protecting the river to make sure that future generations can enjoy the beauty of its flora and fauna. This work will give our fish populations the best opportunity to recover and flourish into the future.

# Fisheries Monitoring of the Ribble Catchment

## 5.0 References

- Arevalo, E. et al., 2018. Effect of food shortage and temperature on age 0+salmonids: a contribution to predict the effects of climatechange. *Journal of Fish Biology*, pp. 642-652.
- Armstrong, J. D., Braithwaite, V. A. & Fox, M., 1998. The response of wild Atlantic salmonparr to acute reductions in water flow.. *Journal of Animal Ecology*, Volume 67, pp. 292-297.
- Baldigo, B. P. & Warren, D. R., 2008. Detecting the Response of Fish Assemblages to Stream Restoration: Effects of Different Sampling Designs. *North American Journal of Fisheries Management*, 28(3), pp. 919-934.
- Caissie, D., 2006. The thermal regime of rivers: A review. *Freshwater Biology*, 51(8), pp. 1389-1406.
- Crozier, W. W. & Kennedy, G. J. A., 1994. Application of semi-quantitative electrofishing to juvenile salmonid stock surveys. *Journal of Fish Biology*, 45(1), pp. 159-164.
- Dugdale, S. J., Bergeron, N. E. & St-Hilaire, A., 2013. Temporal variability of thermal refuges and water temperature patterns in an Atlantic salmon river. *Remote Sensing of Environment*, Volume 136, pp. 358-373.
- Elliott, J. M. & Elliott, J. A., 2006. A 35-year study of stock-recruitment relationships in a small population of sea trout: assumptions, implications and limitations for predicting targets. In: G. Harris & N. Milner, eds. *Sea Trout: Biology, Conservation and Management*. Oxford: Blackwell Publishing, pp. 257-278.
- Elliott, J. M. & Elliott, J. A., 2010. Temperature requirements of Atlantic salmon *Salmo salar*, brown trout *Salmo trutta* and Arctic charr *Salvelinus alpinus*: Predicting the effects of climate change. *Journal of Fish Biology*, 77(8), pp. 1793-1817.
- Fredrich, F., Ohmann, S., Curio, B. & Kirschbaum, F., 2003. Spawning migrations of chub in the River Spree, Germany. *Journal of FishBiology*, Volume 63, pp. 710-723.
- Fulton, T. W., 1904. The rate of growth of fishes.. *22nd Annual Report of the Fishery Board of Scotland*, Issue 3, pp. 141-241.
- Gillson, J. P. et al., 2020. Can aspects of the discharge regime associated with juvenile Atlantic salmon (*Salmo salar* L.) and trout (*S. trutta* L.) densities be identified using historical monitoring data from five UK rivers?. *Fisheries Management and Ecology*, Volume 27, pp. 567-579.
- Jacoby, D. & Gollock, M., 2014. *Anguilla anguilla*. *The IUCN Red List of Threatened Species 2014*. [Online] Available at: <https://www.iucnredlist.org/species/60344/45833138> [Accessed 20 November 2019].
- Jensen, L. F. et al., 2008. Local adaptation in brown trout early life-history traits: implications for climate change adaptability. *Proceedings of The Royal Society B*, 275(1653), p. 2859–2868.
- Kennedy, B. P., Nislow, K. H. & Folt, C. L., 2008. Habitat-Mediated foraging limitations Drive Survival Bottlenecks For Juvenile Salmom. *Ecology*, 89(9), pp. 2529-2541.
- Landergren, P., 2004. Factors affecting early migration of sea trout *Salmo trutta* parr to brackish water.. *Fisheries Research*, Volume 67, p. 283–294.
- Le Cren, E. D., 1951. The length – weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology*, Volume 20, pp. 2-19.
- R core Team, 2019. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>. s.l.:s.n.

## Fisheries Monitoring of the Ribble Catchment


- Ricker, W. E., 1951. Chapter 9. Growth in Length and in Weight. In: W. E. Ricker, ed. *Computation and interpretation of the biological statistics of fish populations..* Ottawa : Fisheries Research Board of Canada, pp. 1-382.
- Solomon, D. J. & Lightfoot, G. W., 2008. *The thermal biology of brown trout and Atlantic salmon*, Bristol: Environment Agency.
- Team, R. C., 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria..
- Team, R. C., n.d. *R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.* URL <https://www.R-project.org/>.. s.l.:s.n.
- Thomas, S. M., Griffiths, S. W. & Ormerod, S. J., 2015. Adapting streams for climate change using riparian broadleaf trees and its consequences for stream salmonids. *Freshwater Biology*, 61(1), pp. 64-77.
- Thomas, S. M., Griffiths, S. W. & Ormerod, S. J., 2015. Adapting streams for climate change using riparian broadleaf trees and its consequences for stream salmonids.. *Freshwater Biology*, 60(1), pp. 64-77.
- Warren, D. R., Mineau, M. M., Ward, E. J. & Kraft, C. E., 2010. Relating fish biomass to habitat and chemistry in headwater streams. *Environmental Biology of Fishes*, Volume 88, pp. 51-61.

# Fisheries Monitoring of the Ribble Catchment

## 6.0 Appendices


### 6.1 Appendix A

#### 13 Year Dataset (87 sites): Brown Trout Fry



Appendix A.1 Cumulative brown trout fry densities on the Ribble catchment for 87 sites fish 2008 – 2020 including 5 year moving average.


#### 13 Year Dataset (87 sites): Atlantic Salmon Fry




Appendix A.2 Cumulative Atlantic salmon fry densities on the Ribble catchment for 87 sites fish 2008 – 2020 including 5 year moving average.

# Fisheries Monitoring of the Ribble Catchment

## 6.2 Appendix B




Appendix B.1 Fork length histogram of all brown trout captured on the Calder catchment 2020. Maximum fork length for 0-year trout = 100 mm at time of survey




Appendix B. 2. Fork length histogram of all brown trout captured on the Hodder catchment 2020. Maximum fork length for 0-year trout = 102 mm at time of survey.

## Fisheries Monitoring of the Ribble Catchment




**Appendix B. 3.** Fork length histogram of all brown trout captured on the Mid-Ribble catchment 2020. Maximum fork length for 0-year trout = 102 mm at time of survey.




**Appendix B. 4.** Fork length histogram of all brown trout captured on the Upper-Ribble catchment 2020. Maximum fork length for 0-year trout = 102 mm at time of survey.

## Fisheries Monitoring of the Ribble Catchment




**Appendix B. 5. Fork length histogram of all Atlantic salmon captured on the Hodder catchment 2020. Maximum fork length for 0-year salmon = 102 mm at time of survey**



**Appendix B. 6. Fork length histogram of all Atlantic salmon captured on the Mid-Ribble catchment 2020. Maximum fork length for 0-year salmon = 101 mm at time of survey**

## Fisheries Monitoring of the Ribble Catchment




*Appendix B. 7. Fork length histogram of all Atlantic salmon captured on the Upper-Ribble catchment 2020. Maximum fork length for 0-year salmon = 102 mm at time of survey*


# Fisheries Monitoring of the Ribble Catchment

## 6.3 Appendix C

DUE TO COVID 19 AND TIME RESTRICTION - Q DATA FROM 2020 WITH THE ADDITION OF DATA FROM 2019 HAS BEEN USED TO CREAT THE CALIBRATION FOR THIS YEAR



Appendix C. 1 Brown trout quantitative fry population relationship between semi-quantitative (5 minutes fry capture) and quantitative electric fishing results (Fry per 100 square) that is LN+1 transformed. Fitted linear regression for 0 + salmonids is produced where  $\ln(y + 1) = 0.2779 + 1.1914 \ln(x + 1)$



Appendix C. 2. Atlantic salmon quantitative fry population relationship between semi-quantitative (5 minutes fry capture) and quantitative electric fishing results (Fry per 100 square) that is LN+1 transformed. Fitted linear regression for 0 + salmonids is produced where  $\ln(y + 1) = 0.098 + 1.1587 \ln(x + 1)$