

Angela Graham Bat Consultancy Service Limited

Office 47, Bury Business Centre, Kay Street, Bury, Lancs, BL9 6BU

tel - 0161 763 6171, fax - 0161 761 7854, mob - 07710 184142

e-mail : bat.consultancy@talktalk.net Company no. 7492656



**Bat Survey Report (preliminary day-time survey):
Mill House Farm,
Chipping Lane,
Chaigley,
BB7 3LS**

OS grid reference:
SD 6911 4152

Commissioned-by:
David Pennington

Survey Date: 25/8/22

Report Date: 1/9/22

Table of Contents

1. Summary	page 3
2. Introduction	page 4
3. Bats and the law	page 6
4. Survey	page 7
5. Limitations of the Survey	page 8
6. Findings	page 8
7. Conclusions/Discussion	page 12
8. Recommendations	page 13
9. References	page 14
Appendix 1 - Angela Graham's Experience	page 15
Appendix 2 - Criteria used in assessing risk of roosting (in the absence of obvious evidence of roosting)	page 17
Appendix 3 - Recommendations for further survey work when the findings of the preliminary survey were negative	page 19
Appendix 4 - Examples of available integrated bat boxes	page 20
Appendix 5 - Examples of house sparrow nesting provision	page 23

1. Summary.

(This summary should be read in conjunction with the conclusions and recommendations.)

This essentially open-ended barn is of brick with an asbestos roof. It has a vaulted ceiling, no hay-loft and no ridge-beam.

It is in a rural location of high quality bat feeding-habitat, where at least 7 species of bat are known or likely to occur within 1km.

The walls have a cavity with some bat access available - at internal wall-heads in particular. While pipistrelle bats are the smallest and most common species, so most likely to make use of some of the opportunities available, a variety of species can be found within wall-cavities, including in winter when hibernating.

Being essentially open-ended, and with an unlined roof, the barn will lack the thermal stability of traditional, rubble-filled, stone barn walls, reducing the likelihood of use by bats other than on an essentially casual basis, though this includes in winter for hibernation.

Good Practice Guidelines suggest even low risk buildings should have a bat activity survey (emergence at dusk or to return-to-roost at dawn) when the findings of the initial survey were negative. In this case I have assessed the building as being at low risk but anticipate that a single bat activity survey is unlikely to coincide with roosting i.e. a negative result is likely to be obtained, but this would not confirm absence of roosting at other times.

A more pragmatic approach would be to undertake a precautionary approach to demolition and recreate roosting potential within walls by means of integrated bat boxes in the new build. The latter would accord with the biodiversity objectives of the National Planning Policy Framework.

Accordingly no demolition should take place between the months of November and March inclusive unless the temperature the night before has remained above 8°C. The roof must be removed with appropriate care first.

Demolition should be by operatives who are familiar-with and sympathetic-to the law relating to bats. They must understand that if at any time a bat or droppings that may have come from a bat are found work **must stop immediately**. They have a legal obligation to do this. As far as practicable the feature that was sheltering the bat/s should be replaced. Further advice **must** then be sought from the bat consultant before work continues, even if the bat has flown off or droppings appear old.

There should be an integrated bat box installed in each elevation of the new-build, towards the eaves.

Care should be taken when planning any lighting on the site, to ensure integrated bat boxes and potential bat flight-lines to and from them are appropriately shielded from light-spill.

Incidentally, care should be taken when dismantling the pile of fence-pots, especially in winter when any bat present would be hibernating.

House sparrow nests were noted at wall-heads. Demolition must not take place while birds are actively nesting. Provide sparrow-terracing in association with the new build. The west elevation is best avoided.

2. Introduction


I was asked to assess the importance of this agricultural building to bats as part of the planning process, prior to its proposed demolition. Incidentally I comment on any issues discovered with respect to other protected/priority/invasive species and species of conservation concern.

This is a brick-built, largely open-ended barn with an unlined, asbestos roof:



Front (east)/north and rear/south elevations

It is in a rural location of high quality bat feeding habitat, linking with the River Hodder about 450m to the north-east:



Location of barn indicated by red circle

The pipistrelle bat (2 species: *Pipistrellus pipistrellus* - the common pipistrelle, and *Pipistrellus pygmaeus* - the soprano pipistrelle) is common and widespread in the area.

Roosts of these species can occur in any building that provides suitable roosting crevices, with the risk of bat presence increased by close proximity to good bat feeding habitat and commuting routes; for example tree-lines, hedges, woodland, scrub and water courses and bodies. The bats use different roosts at different times of year, sometimes singly and sometimes in large groups of females with dependent young. They can move frequently and unpredictably between the roost sites known to them. The majority of house-holders with a roost of this species are unaware of it.

In summer females gather together each with their single off-spring in, sometimes large, maternity colony groups. Disturbance can cause the abandonment of babies (pups). In autumn when the young are independent, females visit males to mate. In winter the bats hibernate and rousing from hibernation - a slow process - can result in a depletion of fat reserves that may compromise the bats' ability to survive the winter. Females become pregnant in spring when their food (insects) becomes available again.

Pipistrelle bats in particular are extremely small, weighing about 5g (the weight of a 2p coin) so need only the smallest of gaps in order to enter to roost; often making use of external features and wall cavities without leaving signs in the loft/interior.

As this is a building with relatively little bat roosting potential, a formal data search hasn't been requested from Lancashire Environmental Records Centre.

A data search from the National Biodiversity Network discovered records of Daubenton's (*Myotis daubentonii*) and Natterer's (*Myotis nattereri*) within 1km, and no additional species within 2kms. The nearest record is of Natterer's about 500m away.

A search of DEFRA's Magic database discovered that bat European Protected Species licences have been granted for common pipistrelle about 1.2kms away and soprano pipistrelle about 1.9kms away.

Other species likely to occur within 1 or 2 kilometres include the brown long-eared (*Plecotus auritus*) - the species most likely to leave evidence of roosting within barns and lofts, the whiskered (*Myotis mystacinus*)/Brandt's (*Myotis brandtii*), which are hard to separate without dna analysis and noctule (*Nyctalus noctula*).

Breeding birds.

Buildings generally can be used by birds of conservation concern (1), such as the house sparrow (*Passer domesticus*), house martin (*Delichon urbicum*), swift (*Apus apus*) and starling (*Sturnus vulgaris*). The house sparrow and starling are both listed in Section 41 of the Natural Environment and Rural Communities Act (NERC) as species "of principal importance for the purpose of conserving biodiversity".

Farm buildings in particular are often used by swallows (*Hirundo rustica*) for nesting and can be used by barn owls (*Tyto alba*). The barn owl has special protection as a "Schedule 1" species. These buildings can be used by other birds of conservation concern such as stock doves (*Columba oenas*) and kestrels (*Falco tinnunculus*) as well as more common species species such as the pied wagtail (*Motacilla alba*) jackdaw (*Corvus*

monedula), wren (*Troglodytes troglodytes*), blue tit (*Cyanistes caeruleus*), great it (*Parus major*), robin (*Erithacus rubecula*) and blackbird (*Turdus merula*).

3. Bats and the Law

All British bats and their roosts are legally protected under the Wildlife and Countryside Act of 1981 (as amended) and the EC Habitats Directive of 1994 as implemented by the Conservation of Habitats and Species Regulations 2017.

Where a development will destroy a bat roost, a Low Impact Class Licence or a European Protected Species Licence (Mitigation Licence) is required before the roost can be interfered with in any way. The former applies in cases where only small numbers of common species of bat are using the building within certain parameters. It usually takes approximately 2 weeks for these licences to be issued, whereas the turn-around time for a full European Protected Species Licence is approximately 7 weeks once the application has been submitted. Any licence issued is a legally binding document.

Licences can only be issued providing planning permission has been granted, where applicable.

When a roost is found, both the bat consultant and the planners have to apply the "three tests" required by Natural England. Essentially these are:

- That the development is necessary for the purpose of “preserving public health or public safety or other imperative reasons of overriding public interest, including those of a social or economic nature and beneficial consequence of primary importance for the environment”;
- That there is “no satisfactory alternative”;
- That the action authorised “will not be detrimental to the maintenance of the population of the species concerned at a favourable conservation status in their natural range”.

N.b. The way in which the necessity is assessed includes whether the client has an alternative that it would be reasonable to expect them to adopt.

Necessary mitigation and compensation measures to ensure the favourable conservation status of bats will be maintained, would include appropriate timing and methodology for the work, including details of how the bats will be provided-for in the long term.

Planners are required by the Government to satisfy themselves before granting planning consent that it would be possible for a licence to be obtained if necessary. Accordingly they are obliged to apply the three tests before issuing consent. For this reason enough survey work has to have been undertaken that the planning authority can evaluate whether or not the three tests can be satisfied and what degree of compensation/enhancement is necessary. To avoid delays in obtaining consent it is in the client's best interest to find out sooner rather than later whether any bat roosting issues need to be addressed.

Natural England, the Government body responsible for administering the law relating to bats, have issued guidelines to planners on how to proceed with respect to bats

Outside the planning system, the onus is on developers/members of the public, to have sufficient investigations undertaken to satisfy themselves (and the authorities in the event of a subsequent investigation), that their actions are unlikely to be in contravention of bat legislation.

N.b. It should always be remembered that bats often roost in places not anticipated by a lay person, such as modern buildings, trees with cavities, and bridges. Some

leave no signs in lofts, as they roost underneath external features such as roof slates, ridges, weather-boarding and cladding.

In the case of a building, tree or other feature not already known to be a bat roost, if bats are found during the course of work, contractors are legally obliged to stop work and seek advice. This should be from an appropriately experienced and licenced bat ecologist.

Breeding birds.

The Wildlife and Countryside Act of 1981 gives protection to the nests of all wild birds whilst being built or in use, including by newly fledged birds that have not left the immediate vicinity of the nest. The bird nesting season is generally considered to be 1st March to 31st July for most species but can extend a number of weeks either side of this depending on the species concerned and weather conditions in that particular year. Natural England cite the nesting season as being 1st March to 31st August.

A consortium of organisations, via their report on "The population status of birds in the UK: Birds of Conservation Concern 5 (2021)" have listed species according to their conservation need based on red, amber, green basis, where red is of the highest conservation concern.

Additional Relevant Legislation and Policy.

Between 1995 and 2010 certain more vulnerable habitats and species were the subject of National or Local Biodiversity Action Plans. This strategy for the protection of biodiversity has been superseded by UK post-2010 Biodiversity Framework, which is largely now implemented at county level. Internationally The Convention on Biodiversity produced a Strategic Plan for Biodiversity 2011-2020. Further to this the EU Biodiversity Strategy was launched in 2011.

Section 41 of the Natural Environment and Rural Communities Act 2006 lists species "of principal importance for the purpose of conserving biodiversity". The list was up-dated in 2014 and includes the brown long-eared bat (*Plecotus auritus*), noctule (*Nyctalus noctula*) soprano pipistrelle (*Pipistrellus pygmaeus*) and 4 other bat species.

The National Planning Policy Framework of 2012 (2) states that "the planning system should contribute to and enhance the natural and local environment" by a number of means, including "minimising impacts on biodiversity and providing net gains in biodiversity where possible, contributing to the Government's commitment to halt the overall decline in biodiversity, including by establishing coherent ecological networks... ."

4. Survey

I made a daytime visit on **25/8/22** to undertake a preliminary survey of the building, assess its likely importance to bats and advise whether or not a precautionary approach or further survey work is needed.

Having been involved with bat survey work for 35 years and consultancy work for 26 years, it is always my objective to carry-out my work in a manner consistent with accepted Good Practice Guidelines (3) and consistent with the code of practice of the CIEEM. I hold Natural England Class Licences CL21 (Annex B) and 18. Amongst other things these cover me to apply for Low Impact Licences for clients and undertake bat

survey work. I also have a CL29 Barn Owl Class Licence. My credentials are expanded-upon in Appendix 1.

As far as possible, I surveyed the building inside and out with the aid of surveyor's ladders, 2 million candle-power torch, camera with 6x optical zoom and binoculars (8x42). Head-torch, 10x 50 binoculars, fibrescope (6 and 13mm heads, extendable to 2m), camera with 18x zoom and mirrors were also available if needed.

I was looking for access to potential roosting places and evidence of their use, such as droppings and staining.

I also take into consideration the surrounding habitat and the range of bat species it appears likely to support, along with the quality of the habitat linkages with the wider area.

The survey was conducted with the needs of different species of bat over the seasons in mind.

Incidentally I comment on any relevant issues discovered with respect to bat feeding habitat and commuting routes, possibly including likely roosting sites nearby, as well as any relevant findings with respect to other protected/invasive species, biodiversity priority species and species of conservation concern.

5. Limitations of the survey

This was a preliminary survey to discover whether there are obvious signs of use by bats and to assess potential for use. If potential exists, usually follow-up work is required at dusk or dawn, possibly at a more appropriate time of year. See Appendix 3.

It should be noted that droppings are the sign most frequently found, but they are often deposited in areas that cannot be easily visualised, if at all, and they can turn to powder quite quickly. They are usually soon washed and blown away from exposed external surfaces so evidence of use often doesn't last long and pipistrelle bats in particular can change roosts frequently. However an assessment has been made of potential bat roosting places associated with the exterior of the building.

As bats often roost in crevices in winter, and are particularly hard to locate when hibernating, the report will highlight any areas that could be used by bats in winter without their presence necessarily being obvious. Bats are particularly vulnerable when hibernating as they cannot rouse quickly to move out of the way.

6. Findings

The building consists of a single, open void:



The roof is unlined and has no ridge beam:



Bat access is easily possible at both gable ends and in places at the ridge.

There are a few gaps into the internal brickwork, where there is clearly a cavity within the wall where bats could roost:



There are a few gaps at beam-insertion-points too, and access to wall-heads:



There was evidence of nesting by birds:



House sparrows use a nearby hedge so it is likely the nests are of house sparrows.

External fascias along the sides of the building are close-fitting and the walls are generally intact, with an occasional minor exception at low level in the west-facing elevation:




There's also a stack of fence-posts alongside the barn currently. Bats have been known to roost/hibernate within stored wood on timber yards.

There is access to wall-heads at the gable ends:



There is some apparent roosting potential available under the asbestos roof where mortar-work is no-longer intact. This was hard to photograph effectively but is illustrated below:



Gap circled in red

There did not appear to be any roosting potential at the northern gable end between vertical planking and the wall or fascia:



7. Conclusions/Discussion

Appendix 2 gives an outline of the criteria used in assessing the level of risk of use by bats.

The walls have a cavity with some bat access available - at internal wall-heads in particular. While pipistrelle bats are the smallest and most common so most likely to make use of some of the opportunities available, a variety of species can be found within wall-cavities, including in winter when hibernating.

The barn is in a high risk location where a range of bat species are likely to occur.

This is an essentially open-ended barn which will lack thermal stability, reducing the likelihood of use by bats other than on an essentially casual basis.

Good Practice Guidelines suggest even low risk buildings should have a bat activity survey (emergence at dusk or to return-to-roost at dawn) when the findings of the initial survey were negative. See Appendix 3.

In this case I have assessed the building as being at low risk but anticipate that a single bat activity survey is unlikely to coincide with roosting i.e. a negative result is likely to be obtained, but this would not confirm absence of roosting at other times.

A more pragmatic approach would be to undertake a precautionary approach to demolition and recreate roosting potential within walls by means of integrated bat boxes in the new build. See Appendix 4. The latter would accord with the biodiversity objectives of the National Planning Policy Framework.

Accordingly demolition should be avoided during the colder winter months in case any bat has chosen to hibernate within a wall; and be carried-out with appropriate care by contractors who are familiar-with and sympathetic-to the law relating to bats.

The bat consultant could attend site to give a tool-box-talk and check wall cavities as far as possible with a fibrescope before they are demolished, though it is anticipated that removal of the roof first will cause sufficient disturbance to deter roosting. This is probably sufficient given that the likelihood of a bat being present at any given time - especially outside winter - is low if not negligible. However, some minor roosting potential was discovered under the asbestos sheeting at wall-heads too.

There should be an integrated bat box in each elevation of the new-build.

Care should be taken when planning any lighting on the site, to ensure any potential roosting features provided are appropriately shielded. (4)

Bear in mind that birds' nests are legally protected throughout the nesting process. As the house sparrow is on the red list of conservation concern, it would also accord with the biodiversity objectives of the National Planning Policy Framework to provide integrated or wall-mounted sparrow-terracing. See Appendix 5.

8. Recommendations

These recommendations should be read in conjunction with the conclusions above.

Do not demolish the walls of the building between the months of November and March inclusive unless the temperature the night before has remained above 8°C.

Demolition should be by operatives who are familiar-with and sympathetic-to the law relating to bats. They must understand that if at any time a bat or droppings that may have come from a bat are found work **must stop immediately**. They have a legal obligation to do this. As far as practicable the feature that was sheltering the bat/s should be replaced. Further advice **must** then be sought from the bat consultant before work continues, even if the bat has flown off or droppings appear old.

The roof must be removed with appropriate care first.

Install an integrated bat box in each elevation of the new build, towards the eaves.

Ensure any external lighting introduced does not illuminate the roosting potential provided, or potential bat flight lines to and from the provision made.

Demolition must not take place while birds are actively nesting.

Provide sparrow-terracing in association with the new build. The west elevation is best avoided.

9. References

1. Eaton, M. A. et al (2015). Birds of Conservation Concern 5: the status of all regularly occurring birds in the UK, Channel Islands and Isle of Man. *British Birds* 114: 723-747.
2. Department for Communities and Local Government (2012). National Planning Policy Framework.
3. Ed. by Collins, J. (2016). Bat Surveys for Professional Ecologists: Good Practice Guidelines - Third Edition. Bat Conservation Trust.
4. Bat Conservation Trust (2018). Guidance Note 8/18: Bats and Artificial Lighting in the UK.

© Angela Graham. September 2022.

Appendix 1 - Angela Graham's Experience.

- I hold Natural England Class Licences CL21 (Annex B) - Registered Consultant 163 - and CL18 (CL18 (2015 11871 - CLS-CLS). CL21 covers me to apply for Low Impact Class Licences for clients - a more stream-lined system for quickly obtaining a licence from Natural England when a roost of a small number of common bat species will be impacted-upon by the development. CL18 covers me for survey/consultancy/scientific work. I have a supplementary licence to possess up to 10 live/dead bat specimens (20123429). I have a CL29 licence to disturb barn owls.
- I'm a member of The Chartered Institute of Ecology and Environmental Management.
- I undertake my work in accordance with the principles outlined in the Bat Conservation Trust's "Good Practice Guidelines".
- I have been involved in bat conservation for over 30 years, initially as a member of the South Lancashire Bat Group from its inception in 1987 and as a volunteer with the Nature Conservancy Council (NCC) - first licenced in 1989. Later, and for many years, I was Co-ordinator/Chair and Trainer for the South Lancashire Bat Group. I trained the people who currently run the group, one of whom has been a Trustee for the Bat Conservation Trust. I was a founder member of the Greater Manchester Bat Group in 2002 and ran the group for 4 years.
- Over the last 27 years I have done increasing numbers of bat surveys on a consultancy basis, firstly part-time then full time from December 2003.
- My experience in applying-for European Protected Species Licences with respect to bats spans over 20 years.
- From 2003 to 2008 I represented the bat groups of the north-west region at national bat worker meetings, hosted by the Bat Conservation Trust.

Other experience includes:

- Attending bat-worker conferences every year since 1988 (mainly England, some in Wales) plus additional symposia on specific topics such as mitigation and woodland bats.
- Helping with winter surveys of underground hibernation sites in Clwyd and north Lancashire.
- Participating in "Bat Detector Workshops" during the 1990s in different areas of the country, concerned with locating bat roosts and feeding sites/commuting routes.
- Sitting on local council "Wildlife Advisory Groups" (WAGs) in the Greater Manchester area from the early 1990s until around 2005.
- Helping local authorities and the Greater Manchester Ecology Unit formulate their Biodiversity Action Plans for bats.
- Administering the bat casework for English Nature (now Natural England) in the South Lancashire and Greater Manchester areas over 1998-2000.
- Assisting with research involving mist netting, harp trapping and radio-tracking.
- Continuing to attend courses run by recognised experts to ensure I stay up-to date both with respect to bat survey-work and conservation, and issues such as health and safety.
- Re-passing the Construction Site (CITB) Operatives test in June 2017.
- Contributing to the Bat Conservation Trust's survey standards guidelines.

Other ecological experience includes:

- Bird watching for fun since 1982 with a general interest in wildlife, ecology and conservation for a similar period.
- Attending short courses and field training with respect to grasses, flowering plants, British mammals including water voles, reptiles and amphibians, non-native invasive plant species, Extended Phase 1 Habitat Surveying, National Vegetation Classification, Environmental Impact Assessment and use of GIS.
- Taking part in British Trust for Ornithology breeding bird surveys annually.
- A year-long sandwich placement assisting with badger research, including radio-tracking.
- Short periods of voluntary work with the Lancashire Wildlife Trust and Royal Society for Protection of Birds.

Appendix 2 - Personally-devised criteria used in assessing risk of roosting (in the absence of obvious evidence at the preliminary survey).

Risk of roosting	Definition	Suggested Action
Nil	Whole of structure/tree can be seen well enough to be sure there are no roosting opportunities.	No need to consider bats further unless development is delayed and potential roosting places might develop in time.
Minimal/negligible	All or most of structure/tree can be seen well enough to suggest there are few, if any, places suitable for roosting and the location does not provide easy access to potential feeding grounds.	Although roosting is thought to be unlikely and therefore the development is unlikely to impact on the favourable conservation status of bats, a precautionary approach should be taken in relevant areas at the time of the work. Further survey work needed only if development delayed.
Low	Whole of structure/tree can be seen well enough to know there are no more than a few openings that could be used by an individual bat or two and/or these provide access to the sorts of features that are likely to be suboptimal due to materials and/or conditions within (eg unstable temperature); and/or the location provides limited access to potential feeding grounds.	Although regular roosting is thought to be relatively unlikely and the development is unlikely to impact on the favourable conservation status of bats, a single survey at dusk or dawn in favourable weather conditions would be appropriate to accord with good practice. This would reduce the extent to which the judgement is based on speculation. If the findings were ambiguous e.g. possible bat emergence and/or considerable bat activity around the building, the survey would need repeating.
Moderate/medium	A small number of openings are present in an area of reasonable habitat, and at least some seem likely to provide access to good conditions for roosting bats, and/or a loft/hay-loft is present that appears to have good qualities for roosting but there were limitations to access or no evidence of bats was found at the time. Cellars may be assessed as potentially being suitable for hibernation in winter,	Further work is needed to better assess the abundance of bat activity in the vicinity and whether or not bats seem to make use of the roosting potential available. To accord with good practice a dusk emergence survey and a dawn return-to-roost survey will be necessary. A second inspection of the interior may also be necessary - if the survey was undertaken in winter for example. As the absence of bats on two occasions wouldn't guarantee absence at other

	but the conditions and/or location aren't optimal.	times, possibly including winter, some precautions would be needed at the time of the work and some roosting potential should be retained/re-created. In the case of cellars and equivalent, inspection in winter is necessary. Some work, for example pointing old stone walls, should be avoided in winter.
High	There is at least one feature that is typical of those favoured by bats for regular roosting and it/they provide access to abundant insect food on-site and/or via good links with the wider natural environment. The feature/s could be suitable for use by a maternity colony, either as a main or satellite roost, or by a territorial male in autumn in the case of pipistrelles, or by individuals or small numbers of bats at any time of year, including winter when hibernating.	The extent to which bats of different species make use of the potential available needs to be investigated by carrying-out at least 3 surveys at dusk and/or dawn spaced over the months of May to September inclusive, possibly extending into April or October if weather conditions are favourable. (Air temperature above 8°C and not more than light rain and/or gentle breeze. I generally plan to do surveys only when the forecast is for 10°C or above.) Maternity colonies have largely disbanded by September, but territorial male pipistrelles may be missed without a survey in September and a lot of smaller roosts are discovered at this time of year. As bats could hibernate unseen in winter and/or roost at other times not covered by the survey work, appropriate precautions would be needed at the time of the work along with maintenance of appropriate potential roosting places.
High - hibernation only	Cave-like places with stable conditions and high humidity, such as cellars can be used for hibernation in winter.	High-risk potential hibernation sites need at least 3 inspections spaced over the winter months as bats will move between sites depending on the weather conditions.

Appendix 3 - Recommendations for further survey work when the findings of the preliminary survey were negative.

Bat Conservation Trust

Table 7.3 Recommended minimum number of survey visits for presence/absence surveys to give confidence in a negative result for structures (also recommended for trees but unlikely to give confidence in a negative result).

Low roost suitability	Moderate roost suitability	High roost suitability
One survey visit. One dusk emergence or dawn re-entry survey ^a (structures). No further surveys required (trees).	Two separate survey visits. One dusk emergence and a separate dawn re-entry survey. ^b	Three separate survey visits. At least one dusk emergence and a separate dawn re-entry survey. The third visit could be either dusk or dawn. ^b

^a Structures that have been categorised as low potential can be problematic and the number of surveys required should be judged on a case-by-case basis (see Section 5.2.9). If there is a possibility that quiet calling, late-emerging species are present then a dawn survey may be more appropriate providing weather conditions are suitable. In some cases, more than one survey may be needed, particularly where there are several buildings in the category.

^b Multiple survey visits should be spread out to sample as much of the recommended survey period (see Table 7.1) as possible; it is recommended surveys are spaced at least two weeks apart, preferably more. A dawn survey immediately after a dusk one is considered only one visit.

Taken from "Bat Surveys for Professional Ecologists: Good Practice Guidelines", 3rd Edition (2)

Table 7.1 Recommended timings for presence/absence surveys to give confidence in a negative result for structures (also recommended for trees but unlikely to give confidence in a negative result).

Low roost suitability	Moderate roost suitability	High roost suitability
May to August (structures) No further surveys required (trees)	May to September ^a with at least one of surveys between May and August ^b	May to September ^a with at least two of surveys between May and August ^b

September surveys are both weather- and location-dependent. Conditions may become more unsuitable in these months, particularly in more northerly latitudes, which may reduce the length of the survey season.

Multiple survey visits should be spread out to sample as much of the recommended survey period as possible; it is recommended that surveys are spaced at least two weeks apart, preferably more, unless there are specific ecological reasons for the surveys to be closer together (for example, a more accurate count of a maternity colony is required but it is likely that the colony will soon disperse). If there is potential for a maternity colony then consideration should be given to detectability. A survey on 31 August followed by a mid-September survey is unlikely to pick up a maternity colony. An ecologist should use their professional judgement to design the most appropriate survey regime.

Taken from "Bat Surveys for Professional Ecologists: Good Practice Guidelines", 3rd Edition (2)

Appendix 4 – examples of available integrated bat boxes

N.b. An internet search for "integrated bat boxes" will bring up types and suppliers of these boxes but advice from the bat consultant should be sought before they are ordered.

EcoSurv Habibat

<http://www.habibat.co.uk/category/bat-boxes>

“Designed to be built into an exterior wall and is available in a variety of faces to match the building. Standard facings of red or blue brick - ideal for new builds - are normally available from stock, or boxes can be made to your specific requirements with a face of brick, stone, timber, or plain (for rendering). Supplied un-pointed.”



Example of Habibat boxes Can also be faced with stone.

Ibstock Ecozone

<http://www.ibstock.com/sustainability-ecozone.asp>





Above: typical unit in situ. Photo © Angela Graham

Green and Blue Bat Block/Brick


<https://www.greenandblue.co.uk/products/bat-block-bat-brick>



Green and Blue Bat Block/Brick in situ

Schwegler 1FR/2FR

An internet search for Schwegler bat boxes will readily bring up suppliers of these boxes



Schwegler 1FR/2FR

Forticrete

Available from building-supply merchants



Above and below - standard colours available. Others available to order



Example of Forticrete boxes in situ

A similar box is available from Habibat. See above listing.

Appendix 5 – examples of house sparrow nesting provision

Sparrow bricks:



Habibat sparrow bricks
EcoSurv Habibat: <http://www.ecosurv.co.uk>



examples of wooden sparrow terracing above and below. An internet search for sparrow terracing or sparrow bricks will provide additional suppliers

Birds

installation examples ▶



Sparrow Terrace 1SP



House Sparrow (Passer domesticus):
These birds thrive in areas cultivated by man, from villages to large cities.

Tree Sparrow (Passer montanus):
Also widely distributed throughout towns and villages as well as the surrounding fields, farms, woodlands and hedges, and woodlands bordering rivers and streams.

General details:
While nesting and raising their young both types of sparrow feed exclusively on insects, especially arthropods (insects with segmented legs). Therefore, their traditional role in keeping down pests and maintaining the balance of nature is a very important one, though not always recognised. Recently the numbers of both species have declined substantially and in some areas they are now rarely seen. Long term studies have confirmed this drastic reduction in their numbers across Europe. The causes include the clearance and monotonous nature of rural areas, the sterility of many

chemicals in agriculture and gardens. The survival of these species is particularly threatened by widespread building renovation and clearance which denies them many nesting possibilities.

Occupants:
House and Tree Sparrows, and in some instances other birds which use nest boxes such as Tits, Redstart and Spotted Flycatcher.

Material:
SCHWEGLER wood-concrete.


Siting:
On all types of houses in built-up areas, and on industrial and agricultural buildings such as barns, sheds and factories.

Installation height:
2 metres or more above ground level.

Cleaning:
Cleaning is advised but is not essential. By turning the safety catch by 180° the front panel can be removed.

Dimensions of brood chambers:
Each: height 16 x width 10.5 x depth approx. 15 cm

External dimensions:
height 24.5 x width 43 x



Installation ▼

Installation Method 1

Simple surface installation using the plugs and screws supplied



▲ Built into insulation of walls

Installation Method 2

Complete installation as a nesting block within brick or concrete walls

To avoid heat-conduction please also use wall insulation, or install at a suitable depth in the wall

